arXiv:1502.05570v1 [math.CA] 19 Feb 2015

Entropies and the derivatives of some Heun functions

Ioan Raşa

Abstract

This short note contains a list of new results concerning the Rényi entropy, the Tsallis entropy, and the Heun functions associated with positive linear operators.

MSC: 33E30, 33C05, 41A36, 94A17

Keywords: Entropy, Heun function, hypergeometric function, positive linear operator.

1 Introduction

This short note contains a list of new results supplementing the articles [10], [11], [12].

The Rényi entropy and the Tsallis entropy associated with positive linear operators have been investigated in [11] and [12]. Section 2 is concerned with two new examples in this direction.

The entropies are naturally related to some Heun functions, as explained in the mentioned articles. Using results from [4] and [13], the derivatives of these Heun functions are studied in Sections 3 and 4.

Detailed proofs will be presented in a forthcoming paper.

2 Integral operators

Let $B_{m-1}\left(x_0, x_1, \dots x_m; \cdot\right)$ be the B-spline function of degree $m-1$ associated to the equidistant points $x_0 < x_1 < \cdots < x_m$. Consider a given function $\sigma \in \mathscr{C}(\mathbb{R})$ such that $\sigma(x) > 0$, $x \in \mathbb{R}$. For $x, t \in \mathbb{R}$, let

$$W_n(x,t) := B_{n-1}\left(x-\sigma(x), x-\sigma(x)+\frac{2\sigma(x)}{n}, \dots, x+\sigma(x); t\right), \quad n \geq 1.$$

The operator $L_n : \mathscr{C}(\mathbb{R}) \to \mathscr{C}(\mathbb{R})$ will be defined by

$$L_n f(x) := \int_{\mathbb{R}} W_n(x,t) f(t) dt, \quad n \geq 1, f \in \mathscr{C}(\mathbb{R}), x \in \mathbb{R}. \tag{1}$$

Several properties of operators of this form are investigated, e.g., in [1], [2], [9].

The Rényi entropy and the Tsallis entropy associated with L_n are, respectively, $-\log \int_{\mathbb{R}} W_n^2(x,t)dt$ and $1-\int_{\mathbb{R}} W_n^2(x,t)dt$; see, e.g., [11] and the references therein. For the operator described by (1), we have

$$\int_{\mathbb{R}} W_n^2(x,t)dt = \frac{c_n}{\sigma(x)}, \quad x \in \mathbb{R},$$

where $c_n > 0$ is a constant depending only on n; in particular $c_1 = \frac{1}{2}$, $c_2 = \frac{2}{3}$, $c_3 = \frac{33}{40}$.

Let $e_i(x) = x^i$, $i = 0, 1, 2$. The variance associated with L_n is defined by $V_n(x) = L_n e_2(x) - (L_n e_1(x))^2$, $x \in \mathbb{R}$; see [11]. In our case,

$$V_n(x) = \frac{\sigma^2(x)}{3n}, \quad x \in \mathbb{R}.$$

Remark that the variance, the Rényi entropy and the Tsallis entropy are synchronous functions of x.

Let now $b_{n,j}(x) := \binom{n}{j} x^j (1-x)^{n-j}$, $j = 0, 1, \dots, n$, $x \in [0,1]$. Then $B_n : \mathscr{C}[0,1] \to \mathscr{C}[0,1]$, $B_n f(x) := \sum_{j=0}^{n} f\left(\frac{j}{n}\right) b_{n,j}(x)$, are the classical Bernstein operators.

The Kantorovich modifications of B_n are defined by

$$Q_n^{[k]} f := \frac{n^k (n-k)!}{n!} D^k B_n f^{(-k)}, \quad f \in \mathscr{C}[0,1], n \geq k \geq 0,$$

where D is the differentiation operator and $f^{(-k)}$ is an antiderivative of order k of f; see [3] and the references therein.

It can be proved that

$$Q_n^{[k]} f = \sum_{j=0}^{n-k} b_{n-k,j} \int_0^1 f(t) B_{k-1}\left(\frac{j}{n}, \frac{j+1}{n}, \dots, \frac{j+k}{n}; t\right) dt.$$

Therefore, in order to compute the Rényi entropy and the Tsallis entropy associated with $Q_n^{[k]}$, we need

$$S_n^{[k]}(x) := \int_0^1 \left(\sum_{j=0}^{n-k} b_{n-k,j}(x) B_{k-1}\left(\frac{j}{n}, \frac{j+1}{n}, \dots, \frac{j+k}{n}; t\right) \right)^2 dt.$$

In fact, it can be proved that

$$\begin{aligned} S_{n+2}^{[2]}(x) &= \frac{n+2}{3(n+1)4^n} \sum_{i=0}^{n} (3n-2i+2) 4^i \binom{2i}{i} \binom{2n-2i}{n-i} \left(x - \frac{1}{2}\right)^{2i} \\ &= \frac{n+2}{3\pi} \int_0^{\pi} \left(1 - 4x(1-x)\sin^2\frac{\phi}{2}\right)^n \left(1 + 2\cos^2\frac{\phi}{2}\right) d\phi. \qquad (2) \end{aligned}$$

The variance associated with $Q_{n+2}^{[2]}$ is

$$V_{n+2}^{[2]}(x) = \frac{n}{(n+2)^2}x(1-x) + \frac{1}{6(n+2)^2}, \quad x \in [0,1].$$

Again it follows that the variance, the Rényi entropy and the Tsallis entropy associated with $Q_{n+2}^{[2]}$ are synchronous functions.

3 Heun functions and their derivatives

With classical notation for Heun functions $Hl(a,q;\alpha,\beta;\gamma,\delta;x)$ and hypergeometric functions ${}_2F_1(a,b;c;x)$ we have (see [12])

$$\begin{aligned} Hl\left(\frac{1}{2},q;2q,1;1,1;x\right) &= (1-x)^{-2q}{}_2F_1\left(q,q;1;\left(\frac{x}{x-1}\right)^2\right) \\ &= \frac{1}{\pi}\int_0^\pi \left(1-4x(1-x)\sin^2\frac{\phi}{2}\right)^{-q} d\phi. \end{aligned} \tag{3}$$

The integral representation can be compared with (2).
Combined with

$${}_2F_1(a,b;c;x) = (1-x)^{-a}{}_2F_1\left(a,c-b;c;\frac{x}{x-1}\right),$$

(3) leads to

$$Hl\left(\frac{1}{2},q;2q,1;1,1;x\right) = (1-2x)^{-q}{}_2F_1\left(q,1-q;1;\frac{x^2}{2x-1}\right). \tag{4}$$

With notation from [10], [12], let

$$F_n(x) := \sum_{k=0}^{n}\left(\binom{n}{k}x^k(1-x)^{n-k}\right)^2,$$

$$G_n(x) := \sum_{k=0}^{\infty}\left(\binom{n+k-1}{k}x^k(1+x)^{-n-k}\right)^2,$$

$$U_n(x) := \sum_{k=0}^{n}\left(\binom{n}{k}x^k(1+x)^{-n}\right)^2,$$

$$J_n(x) := \sum_{k=0}^{\infty}\left(\binom{n+k}{k}x^k(1-x)^{n+1}\right)^2.$$

It was proved in [12] that

$$F_n(x) = Hl\left(\frac{1}{2},-n;-2n,1;1,1;x\right), \tag{5}$$

$$G_n(x) = Hl\left(\frac{1}{2}, n; 2n, 1; 1, 1; -x\right), \tag{6}$$

and, moreover, the polynomial Heun function $F_n(x)$ and the rational Heun function $G_n(-x)$ are related by

$$G_n(-x) = (1-2x)^{1-2n} F_{n-1}(x). \tag{7}$$

Similarly, $U_n(x)$ and $J_n(x)$ can be expressed as

$$U_n(x) = F_n\left(\frac{x}{x+1}\right), \tag{8}$$

$$J_n(x) = \left(\frac{1-x}{1+x}\right)^{2n+1} F_n\left(\frac{1}{1-x}\right). \tag{9}$$

Recall that the Legendre polynomials $P_n(x)$ are related to ${}_2F_1$ by

$${}_2F_1(-n, n+1; 1; x) = P_n(1-2x). \tag{10}$$

From (5), (4) and (10) we get

$$F_n(x) = (1-2x)^n P_n\left(\frac{2x^2-2x+1}{1-2x}\right). \tag{11}$$

This formula was proved (with a different method) by Thorsten Neuschel [7] and Geno Nikolov [8]; it was used in order to prove a conjecture involving the polynomials $F_n(x)$.

The derivative of a Heun function $Hl(a, q; \alpha, \beta; \gamma, \delta; x)$ satisfying

$$q = a\alpha\beta \tag{12}$$

was studied in [4]. From the corresponding results we infer, for $\gamma \neq 0, -1, -2, \ldots$,

$$\begin{aligned} &\frac{d}{dx} Hl\left(\frac{1}{2}, \frac{1}{2}\alpha\beta; \alpha, \beta; \gamma, \gamma; x\right) \\ = \quad &\frac{\alpha\beta}{\gamma}(1-2x) Hl\left(\frac{1}{2}, \frac{1}{2}(\alpha+2)(\beta+2); \alpha+2, \beta+2; \gamma+1, \gamma+1; x\right), \end{aligned} \tag{13}$$

$$\begin{aligned} &\frac{d}{dx} Hl\left(\frac{1}{2}, \frac{1}{2}\alpha\beta; \alpha, \beta; \gamma, \gamma; x\right) \\ = \quad &\frac{\alpha\beta}{\gamma}(1-2x)^{2\gamma-\alpha-\beta-1} Hl\left(\frac{1}{2}, \frac{1}{2}(2\gamma-\alpha)(2\gamma-\beta); 2\gamma-\alpha, 2\gamma-\beta; \gamma+1, \gamma+1; x\right). \end{aligned} \tag{14}$$

The equality between the right-hand sides of (13) and (14) follows also from line 3 in Table 2 of [6]. Let us remark that the Heun functions in these right-hand sides satisfy also the condition (12), so that it is possible to express their derivatives in terms of other Heun functions.

From (5) and (13) we get

$$\frac{d}{dx}F_n(x) = 2n(2x-1)Hl\left(\frac{1}{2}, 3-3n; 2-2n, 3; 2, 2; x\right), \tag{15}$$

and finally, for $i = 0, 1, \ldots, n$,

$$Hl\left(\frac{1}{2}, (i-n)(2i+1); 2(i-n), 2i+1; i+1, i+1; x\right) \tag{16}$$

$$= \frac{(2i)!!}{(2i-1)!!}4^{-n}\binom{n}{i}^{-1}\sum_{j=0}^{n-i}4^j\binom{i+j}{i}\binom{2i+2j}{i+j}\binom{2n-2i-2j}{n-i-j}\left(x-\frac{1}{2}\right)^{2j}.$$

To conclude this section, let us remark that the Heun function from (3) satisfies the condition (33) from [5]; the consequence of this fact will be investigated elsewhere.

4 A confluent Heun function

Let $u(x) = HC(p, \gamma, \delta, \alpha, \sigma; x)$ be the solution of the confluent Heun equation

$$u''(x) + \left(4p + \frac{\gamma}{x} + \frac{\delta}{x-1}\right)u'(x) + \frac{4p\alpha x - \sigma}{x(x-1)}u(x) = 0, \tag{17}$$

with $u(0) = 1$ (See [13]). From [13, (21)] we get

$$\frac{d}{dx}HC(p, \gamma, 0, \alpha, 4p\alpha; x) = -\frac{\sigma}{\gamma}HC(p, \gamma+1, 0, \alpha+1, 4p(\alpha+1); x), \tag{18}$$

$$\frac{d}{dx}HC(p, \gamma, 0, \alpha, 4p\alpha; x) = \frac{\sigma}{\gamma}(x-1)HC(p, \gamma+1, 2, \alpha+2, 4p(\alpha+1)-\gamma-1; x). \tag{19}$$

With notation from [10], [12], let

$$K_n(x) := \sum_{k=0}^{\infty}\left(e^{-nx}\frac{(nx)^k}{k!}\right)^2.$$

Then (see [10], [12]),

$$xK_n''(x) + (4nx+1)K_n'(x) + 2nK_n(x) = 0. \tag{20}$$

We get immediately

$$K_n(x) = HC\left(n, 1, 0, \frac{1}{2}, 2n; x\right). \tag{21}$$

From (19) and (21) it follows that

$$HC\left(n, 2, 2, \frac{5}{2}, 6n-2; x\right) = \frac{1}{2n(x-1)}K_n'(x). \tag{22}$$

Similarly, from (18) and (21),

$$HC\left(n, 2, 0, \frac{3}{2}, 6n; x\right) = -\frac{1}{2n} K_n'(x). \tag{23}$$

Now applying repeatedly (18) we can generalize (23) to

$$HC\left(n, j+1, 0, \frac{2j+1}{2}, 2n(2j+1); x\right) = \frac{K_n^{(j)}(x)}{K_n^{(j)}(0)}, \quad j \geq 0, \tag{24}$$

where

$$K_n^{(j)}(0) = (-2n)^j \sum_{i=0}^{[j/2]} \binom{j}{2i} \binom{2i}{i} 4^{-i}. \tag{25}$$

From (20) we get

$$\left(K_n^{(j)}\right)'' + \left(4n + \frac{j+1}{x}\right) \left(K_n^{(j)}\right)' + \frac{2n(2j+1)}{x} K_n^{(j)} = 0, \tag{26}$$

and this provides an alternative proof of (24).

References

[1] M. Campiti, I. Raşa, C. Tacelli, *Steklov operators and their associated semigroups*, Acta Sci. Math. (Szeged) 74 (2008), 171-189.

[2] M. Campiti, I. Raşa, C. Tacelli, *Steklov operators and semigroups in weighted spaces of continuous real functions*, Acta Math. Hungar. 120 (2008), 103-125.

[3] H. Gonska, M. Heilmann, I. Raşa, *Kantorovich operators of order k*, Numer. Funct. Anal. Optimiz. 32 (2011), 717-738.

[4] A. Ishkhanyan, K.A. Suominen, *New solutions of Heun's general equation*, J. Phys. A: Math. Gen. 36 (2003), L81-L85.

[5] T.A. Ishkhanyan, T.A. Shahverdyan, A.M. Ishkhanyan, *Hypergeometric expansions of the solutions of the general Heun equation governed by two-term recurrence relations for expansions coefficients*, arXiv:1403.7863.

[6] R.S. Maier, *On reducing the Heun equation to the hypergeometric equation*, J. Differential Equations 213 (2005), 171-203.

[7] Th. Neuschel, Unpublished manuscript, Univ. of Trier, 2 pp, 2012.

[8] G. Nikolov, *Inequalities for ultraspherical polynomials. Proof of a conjecture of I. Raşa*, J. Math. Anal. Appl. 418 (2014) 852-860.

[9] D. Popa, I. Raşa, *Steklov averages as positive linear operators*, to appear in FILOMAT.

[10] I. Raşa, *Special functions associated with positive linear operators*, arXiv:1409.1015v2 [math.CA].

[11] I. Raşa, *Rényi entropy and Tsallis entropy associated with positive linear operators*, arXiv:1412.4971v1 [math.CA].

[12] I. Raşa, *Entropies and Heun functions associated with positive linear operators*, Preprint 2015.

[13] V.A. Shahnazaryan, T.A. Ishkhanyan, T.A. Shahverdyan, A.M. Ishkhanyan, *New relations for the derivative of the confluent Heun function*, Armenian J. Phys. 5 (2012), 146-156.